

A Novel Bunyavirus-Like Virus of Trypanosomatid Protist Parasites

Natalia S. Akopyants,^a Lon-Fye Lye,^a Deborah E. Dobson,^a Julius Lukeš,^{b,c}  Stephen M. Beverley^a

Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, Missouri, USA^a; Institute of Parasitology, Biology Centre, and Faculty of Sciences, University of South Bohemia, České Budějovice, Czech Republic^b; Canadian Institute for Advanced Research, Toronto, Ontario, Canada^c

We report here the sequences for all three segments of a novel RNA virus (LepmorLBV1) from the insect trypanosomatid parasite *Leptomonas moramango*. This virus belongs to a newly discovered group of bunyavirus-like elements termed Leishbunyaviruses (LBV), the first discovered from protists related to arboviruses infecting humans.

Received 30 May 2016 Accepted 17 June 2016 Published 4 August 2016

Citation Akopyants NS, Lye L-F, Dobson DE, Lukeš J, Beverley SM. 2016. A novel bunyavirus-like virus of trypanosomatid protist parasites. *Genome Announc* 4(4):e00715-16. doi:10.1128/genomeA.00715-16.

Copyright © 2016 Akopyants et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Stephen M. Beverley, beverley@wusm.wustl.edu.

Bunyaviridae comprise >350 species, often arthropod-borne and including serious human, animal, and plant pathogens (1). In searching for novel viruses from trypanosomatid protists (Kinetoplastida, supergroup Excavata), we found a new viral element within the insect parasite *Leptomonas moramango* (2).

RNA was prepared from an uncloned strain of parasites (2) using TRIzol reagent (Thermo Fisher), treatment with DNase I (Thermo Fisher), and purified with RNA Clean & Concentrator-25 (Zymo Research). Replicative viral double-stranded RNAs (dsRNAs) were visualized following treatment with S1 nuclease (Thermo Fisher), separation by agarose gel electrophoresis, and staining by ethidium bromide (3). Three prominent bands with sizes of about 6, 1.3, and 0.7 kb were observed (L, M, and S, respectively). Total RNA was depleted of rRNA with the RiboZero kit (Illumina), fragmented to 200 to 600 nucleotides (nt), and used as a template for the generation of multiplexed TruSeq cDNA libraries. These were sequenced (2 × 101 cycles, paired-end reads) on the HiSeq 2500 (Illumina). A total of 45,411,994 reads were obtained, of which 722,520 (1.6%) were assigned to viral segments (CLC Genomics Workbench).

For each segment, we identified two related similarly sized contigs (L, 5,982 and 6,029 nt; M, 1,205 and 1,154 nt; S, 662 and 820 nt). Within each pair, one was more abundant, as judged by read depth coverage (L, 1,634 or 471 nt; M, 12,453 or 225 nt; S, 53,243 or 622 nt); we arbitrarily grouped the more abundant segments as “a” and refer to the viruses as LepmorLBV1a and 1b. The G+C compositions of LepmorLBV1s were similar to each other and to that of the meta-transcriptomic assembly (38 to 41%). We were able to detect 9-nt “panhandle” repeats (4, 5) on 6/12 termini within the metatranscriptomic contigs.

Comparisons between the LepmorLBV1a/b segment open reading frames (ORFs) showed significant nucleotide and amino acid identity (52 to 54% and 37 to 50%, respectively). L segments displayed a single ORF encoding 1,979 or 1,982 amino acids (aa), bearing motifs found typically in bunyavirus RNA-dependent RNA polymerases (RdRp) (6, 7). M segments encoded a single ORF of 322 or 321 aa; no database hits were found, but the protein(s) were predicted to have a hydrophobic N-terminal signal sequence. S segments predicted proteins of 165 and 168 aa, which

showed structural similarity to the nucleocapsid of other bunyaviruses (4, 5, 8).

Phylogenetic analysis of the RdRp domain showed that LepmorLBV1s grouped together with bunyaviruses, relative to other viral outgroups, but as a clearly distinct clade well separated from *Phlebovirus* or other genera. Additional affinities to *Bunyaviridae* include a negative single-stranded RNA genome, terminal repeated panhandle sequences important for replication and transcription, and three segments typically encoding the viral RdRp, envelope glycoproteins, and the nucleocapsid (4, 5, 7). The *L. moramango* virus thus resembles a group of related viruses discovered recently in the closely related human parasite *Leishmania*, which we have termed *Leishbunyavirus* (LBV).

L. moramango may bear cytoplasmic virus-like particles (2) that could correspond to LepmorLBV1. Future studies will establish the evolutionary distribution of LBVs and their potential role in host pathogenicity (9).

Accession number(s). The sequences of all segments of LepmorLBV1a and 1b were deposited in GenBank under the accession numbers [KX280012](https://www.ncbi.nlm.nih.gov/nuclink/KX280012) to [KX280017](https://www.ncbi.nlm.nih.gov/nuclink/KX280017).

ACKNOWLEDGMENTS

We thank members of our laboratories, Vyacheslav Yurchenko, and Nicolas Fasel for discussions.

This work was supported by NIH AID R56 AI099364 and AI29646 to S.M.B. Next-generation sequencing was performed at the Washington University School of Medicine, Department of Genetics Genome Technology Access Center (partially supported by grants NCI Cancer Center Support P30 CA91842 and NCRR ICTS/CTSA UL1 TR000448).

FUNDING INFORMATION

This work, including the efforts of Natalia S. Akopyants, Lon-Fye Lye, Deborah E. Dobson, and Stephen M. Beverley, was funded by HHS | NIH | National Cancer Institute (NCI) (NCI P30 CA91842). This work, including the efforts of Natalia S. Akopyants, Lon-Fye Lye, Deborah E. Dobson, and Stephen M. Beverley, was funded by HHS | National Institutes of Health (NIH) (R01-AI29646). This work, including the efforts of Natalia S. Akopyants, Lon-Fye Lye, Deborah E. Dobson, and Stephen M. Beverley, was funded by HHS |

NIH | National Center for Research Resources (NCRR) (UL1 TR000448).

REFERENCES

1. Plyusnin A, Beaty BJ, Elliott RM, Goldbach R, Kormelink R, Lundkvist KA, Schmaljohn CS, Tesh RB. 2012. Family *Bunyaviridae*, p 725–741. In King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ (ed), *Virus taxonomy: classification and nomenclature of viruses: ninth report of the International Committee on Taxonomy of Viruses*. Elsevier Academic, San Diego, CA.
2. Yurchenko V, Votýpka J, Tesařová M, Klepetková H, Kraeva N, Jirků M, Lukeš J. 2014. Ultrastructure and molecular phylogeny of four new species of monoxenous trypanosomatids from flies (Diptera: Brachycera) with redefinition of the genus *Wallaceina*. *Folia Parasitol (Praha)* 61:97–112.
3. Beiting DP, Peixoto L, Akopyants NS, Beverley SM, Wherry EJ, Christian DA, Hunter CA, Brodsky IE, Roos DS. 2014. Differential induction of TLR3-dependent innate immune signaling by closely related parasite species. *PLoS One* 9:e88398. <http://dx.doi.org/10.1371/journal.pone.0088398>.
4. Guu TS, Zheng W, Tao YJ. 2012. *Bunyavirus*: structure and replication. *Adv Exp Med Biol* 726:245–266. http://dx.doi.org/10.1007/978-1-4614-0980-9_11.
5. Walter CT, Barr JN. 2011. Recent advances in the molecular and cellular biology of bunyaviruses. *J Gen Virol* 92:2467–2484. <http://dx.doi.org/10.1099/vir.0.035105-0>.
6. Reguera J, Weber F, Cusack S. 2010. *Bunyaviridae* RNA polymerases (L-protein) have an N-terminal, influenza-like endonuclease domain, essential for viral *cap*-dependent transcription. *PLoS Pathog* 6:e1001101. <http://dx.doi.org/10.1371/journal.ppat.1001101>.
7. Gerlach P, Malet H, Cusack S, Reguera J. 2015. Structural insights into bunyavirus replication and its regulation by the vRNA promoter. *Cell* 161:1267–1279. <http://dx.doi.org/10.1016/j.cell.2015.05.006>.
8. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. 2015. The Phyre2 Web portal for protein modeling, prediction and analysis. *Nat Protoc* 10:845–858. <http://dx.doi.org/10.1038/nprot.2015.053>.
9. Hartley MA, Drexler S, Ronet C, Beverley SM, Fasel N. 2014. The immunological, environmental, and phylogenetic perpetrators of metastatic leishmaniasis. *Trends Parasitol* 30:412–422. <http://dx.doi.org/10.1016/j.pt.2014.05.006>.